

BioLean II® Supplement Facts

Order from www.MyNaturalSupplements.com

AM Serving Size 2 Brown Tablets, 1 White Tablet

PM Serving Size 1 Brown Tablet

56 Servings Per Container (28 Daily Servings)

Amount Per Serving	AM	PM	% Daily Value*
Calcium (as calcium carbonate, calcium phosphate dibasic)	206.76 mg	91.88 mg	AM=20% PM= 9%
BioLean II Proprietary Blend			+
Caffeine (as Guarana Seed 50% Extract, Yerba Mate Leaf 10% Extract, Green Tea Leaf 40% Extract)	167 mg	83.5 mg	
Citrus Aurantium Fruit 30% Extract (Advantra Z®)			+
Schizandrae berry			+
Gymnema Sylvestre Leaf 25% Extract			+
Rehmannia Root			+
Hawthorne berry			+
Jujube Seed			+
Alisma root			+
Angelicae dahuricae Radix			+
Epemidium grandiflorum Radix			+
Poria Cocos Mushroom			+
Rhubarb Root			+
Angelicae sinensis Radix			+
Codonopsis root			+
Euconium bark			+
Panax notoginseng Radix			+
L-tyrosine	196 mg		+
L-phenylalanine	196 mg		+
L-carnitine (as L-carnitine bitartrate)	8 mg		+

* Percentages are based on a 2,000 calorie/day diet.

+ Daily Value not established.

Other Ingredients (Herbal Blend): Calcium carbonate, starch, stearic acid, cellulose, hydroxypropylcellulose, croscarmellose, sodium, magnesium stearate, silicon dioxide.

Other Ingredients (Amino Acid): Calcium phosphate dibasic, stearic acid, silicon dioxide, croscarmellose sodium, hydroxypropylcellulose, magnesium stearate, ethylcellulose.

Recommended Use: AM Serving - Adults take one white tablet and two brown tablets with low calorie food. PM Serving - Adults may take one brown tablet with low calorie food. If using BioLean® II dietary supplement for the first time, limit daily intake to one white tablet and one brown tablet on days one and two, and one white tablet and two brown tablets on day three. Needs vary with each individual.

Warning: Not for use by children under the age of 18. Do not use if you are pregnant or nursing, or if you have a family history of heart disease, thyroid disease, diabetes, high blood pressure, depression or other psychiatric condition, glaucoma, difficulty urinating, prostate enlargement, or seizure disorder, if you are using a monoamine oxidase inhibitor (MAOI), consult a health professional before using this product. Exceeding recommended serving may cause serious adverse effects. Discontinue use and consult your health professional if dizziness, sleeplessness, severe headache, heart palpitations or other similar symptoms occur. The recommended dose contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, food, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat.

Phenylketonurics: Contains phenylalanine 196mg per AM Serving