

BioLean Free® Supplement Facts

Order from www.MyNaturalSupplements.com

Serving Size 4 tablets * 28 Servings Per Container

Amount Per Serving	% Daily Value*
Niacin (as niacinamide) 40 mg	200%
Vitamin B6 (as pyridoxine HCl) 16 mg	800%
Chromium (as chromium Chelavite® chloride) 400 mcg	332%
Potassium (as potassium citrate) 10 mg	2.8%
Thiamin (as thiamin mononitrate) 1.5 mg	100%
Proprietary Blend Standardized Botanical Extracts 2972 mg	+
Green tea leaf extract (10% methylxanthines)	+
Yerbe maté leaf extract (10% methylxanthines)	+
Uva ursi leaf (20% arbutin)	+
Guarana seed (22% methylxanthines)	+
Quebracho bark extract (10% quebrachine)	+
Proprietary Blend Non-Irradiated Pure Herbs and Thermogenic Spices 1440 mg	+
Gotu kola leaf (Centella asiatica)	+
Ceylon cinnamon bark	+
Chinese horseradish root	+
Jamaican ginger root	+
Turmeric rhizome	+
Nigerian cayenne pepper (fruit)	+
English mustard seed	+
Ho shou wu root	+
Ginkgo biloba leaf (24% ginkgoflavoneslycosides and 6% bilobalides)	+
L-Tyrosine 500 mg	+
L-Methionine 100 mg	+
Vanadium (as BMOV) 400 mcg	+

* Percentages are based on a 2,000 calorie/day diet.

+ Daily Value not established.

Other Ingredients: Dicalcium phosphate, cellulose, cellulose gum, vegetable stearic acid, silica, vegetable magnesium stearate and vegetable resin glaze.

Directions: Adults (18 years and older) may take 4 tablets in the mid to late morning with a low calorie food. Needs may vary with each individual. Some persons may require less than 4 tablets, or may prefer taking 3 tablets mid morning and 1 additional tablet mid afternoon to achieve optimum results. Do not exceed recommended daily amounts. Store in a cool dry place.

Caution: Not for use by children, pregnant women, or lactating women. Consult your physician before using this product if you are taking appetite suppressing drugs or cardiovascular medication. Also consult your physician if you have hypertension, heart disease, arrhythmias, prostatic hypertrophy, glaucoma, liver disease, renal disease, or diabetes. Do not use if you have hyperthyroidism, psychosis, Parkinson's Disease, or are taking Monoamine Oxidase Inhibitors. BioLean Free® should not be taken on the same day as BioLean II®. It is recommended that you minimize your caffeine intake while consuming BioLean Free®. If allergic symptoms develop, discontinue use.